

Stephen K. Aoki, MD

University Orthopedic Center & Primary Children's Medical Center

Osteochondritis Dissecans (Knee)

Surgery:

- 1-2 Hours Long
- Same Day Surgery
- Anesthesia may offer & recommend a nerve block for pain control

Surgical Goals:

- Treatment varies depending on size & stability of the lesion
- Final decision of surgical management is done at time of surgery (drilling vs. re-fixation vs. graft procedure)
 - Decrease Pain/Improve Function

General Timeline After Surgery:

- 0-6 weeks Post-Op: Toe touch weight bearing with crutches. In immobilizer brace while using crutches for the week after surgery. Okay to come out of brace while sitting and lying to work on full range of motion. Okay to perform exercises including: quad sets, straight leg raises, patellar mobilizations & stretching
 - Start formal Physical Therapy 2 weeks after surgery date

(Find a physical therapist that's close to home and covered by insurance)

- 6-9 weeks Post-Op: Begin weaning off of crutches unless otherwise notified. Full range of motion.
 - Okay to add in more stretching and strengthening exercises, but no impact activity
 - 9-12 weeks Post-Op: Full weight bearing without crutches.

Gradually increase to full activities at 3 months post-op as long as x-rays demonstrate proper healing

- New X-Rays are taken at every post-op visit to monitor healing
- Typical Post-Op Appointments: 2 weeks, 3 months, 6 months, and 1 year out from surgery
- **If the OCD lesion requires screw re-fixation, a hardware removal surgery takes place between 6-12 weeks after the original surgery date**

Dr. Aoki's Team Contact Information: FOR MORE INFORMATION VISIT WWW.AOKIMD.COM

Clinic/Pre & Post-Op Questions

Phone: 801-587-2557

Clinic/Appointment Questions

Surgery Scheduling

Phone: 801-587-7487 Email: reed.omdal@utah.com

Primary Children's: 801-662-5607

Email: angel.waddoups@utah.com UUOC: 801-587-7187

Clinic Fax Number: 801-587-3990